arXiv:math-ph/0011004v1 2 Nov 2000

On the time evolution in totally constrained systems with weakly vanishing Hamiltonian[*]

February 7, 2008

S. I. MUSLIH
Dept. of Physics
Al-Azhar University
Gaza, Palestine

Abstract

The Dirac method treatment for finite dimensional singular systems with weakly vanishing Hamiltonin leads to obtain the equation of motion in terms of Parameter τ. To obtain the correct equations of motion one should use gauge fixing of the form $\tau - f(t) = 0$. It is shown that the canonical method leads to describe the evolution in both standard and constrained finite dimensional systems with weakly vanishing Hamiltonian in terms of the physical time t, without using any gauge fixing conditions. Besides the operator quantization of these systems is investigated using the canonical method and it is shown that the evolution of the state Ψ with the time t is described by the Schrödinger equation $i\frac{\partial\Psi}{\partial t} = \hat{H}\Psi$. The extension of this treatment to infinite dimensional systems is given.

---

[*] e-mail: *sami_muslih*@hotmail.com

1 Introduction

The canonical method [1-4] gives the set of Hamilton - Jacobi partial differential equations [HJPDE] as

$$\begin{aligned}
&H_{\alpha}^{'}(t_{\beta}, q_{a}, \frac{\partial S}{\partial q_{a}}, \frac{\partial S}{\partial t_{a}}) = 0, \\
&\alpha, \beta = 0, n-r+1, ..., n, a = 1, ..., n-r,
\end{aligned} \tag{1}$$

where

$$H_{\alpha}^{'} = H_{\alpha}(t_{\beta}, q_{a}, p_{a}) + p_{\alpha}, \tag{2}$$

and H_0 is defined as

$$\begin{aligned}
&H_0 = p_a w_a + p_{\mu} \dot{q_a}|_{p_{\nu}=-H_{\nu}} - L(t, q_i, \dot{q_{\nu}}, \dot{q_a} = w_a), \\
&\mu,\ \nu = n-r+1, ..., n.
\end{aligned} \tag{3}$$

The equations of motion are obtained as total differential equations in many variables as follows:

$$dq_a = \frac{\partial H_{\alpha}^{'}}{\partial p_a} dt_{\alpha}; \tag{4}$$

$$dp_a = -\frac{\partial H_{\alpha}^{'}}{\partial q_a} dt_{\alpha}; \tag{5}$$

$$dp_{\beta} = -\frac{\partial H_{\alpha}^{'}}{\partial t_{\beta}} dt_{\alpha}; \tag{6}$$

$$\begin{aligned}
dZ = &(-H_{\alpha} + p_a \frac{\partial H_{\alpha}^{'}}{\partial p_a}) dt_{\alpha}; \\
&\alpha, \beta = 0, n-r+1, ..., n, a = 1, ..., n-r
\end{aligned} \tag{7}$$

where $Z = S(t_{\alpha}; q_a)$. The set of equations (4-7) is integrable [3,4] if

$$dH_0^{'} = 0, \tag{8}$$

$$dH_{\mu}^{'} = 0, \mu = n-r+1, ..., n. \tag{9}$$

If conditions (8)and (9) are not satisfied identically, one considers them as new constraints and again testes the consistency conditions. Hence, the canonical formulation leads to obtain the set of canonical phase space coordinates q_a and p_a as functions of t_α, besides the canonical action integral is obtained in terms of the canonical coordinates.The Hamiltonians $H_\alpha^{'}$ are considered as the infinitesimal generators of canonical transformations given by parameters t_α respectively.

For the quantization of constrained systems we can use the Dirac's method of quantization [5,6]. In this case we have

$$H_\alpha^{'}\Psi = 0, \quad \alpha = 0, n-r+1, ..., n, \tag{10}$$

where Ψ is the wave function. The consistency conditions are

$$[H_\mu^{\prime}, H_\nu^{\prime}]\Psi = 0, \quad \mu, \nu = 1, ..., r, \tag{11}$$

where$[,]$ is the commutator. The constraints $H_\alpha^{\prime}$ are called first- class constraints if they satisfy

$$[H_\mu^{\prime}, H_\nu^{\prime}] = C_{\mu\nu}^{\gamma} H_\gamma^{\prime}. \tag{12}$$

In the case when the Hamiltonians $H_\mu^{\prime}$ satisfy

$$[H_\mu^{\prime}, H_\nu^{\prime}] = C_{\mu\nu}, \tag{13}$$

with $C_{\mu\nu}$ do not depend on q_i and p_i, then from (11) there arise naturally Dirac' brackets and the canonical quantization will be performed taking Dirac's brackets into commutators.

On the other hand, The path integral quantization is an alternative method to perform the quantization of constrained systems. If the system is integrable then one can solve equations (4-6) to obtain the canonical phase-space coordinates as

$$q_a \equiv q_a(t, t_\mu), \quad p_a \equiv p_a(t, t_\mu), \ \ \mu = 1, ..., r, \tag{14}$$

then we can perform the path integral quantization using Muslih method [7-10] with the action given by (7).

The aim of this paper is to to discuss the meaning of the time evolution in constrained systems with vanishing Hamiltonian.

Even though systems with this property are treated in literature [5,6,11,12], there exists ambiguity in describing the significance of the time. For example some [13] claim that the standard Hamiltonian dynamics expresses the evolution in terms of the clock time t, while the systems with vanishing Hamiltonian describe the evolution in terms of another parameter time τ. The canonical method [1-4] removes this ambiguity since we have more than one Hamiltonian,which leads us to obtain the equations of motion as total differential equations in my variables.

Now we will obtain the equations of motion for four constrained systems with vanishing Hamiltonians and demonstrate the fact that the dynamical variables are obtained in terms of time ($x_0 = t$) without using any gauge fixing conditions.

2 examples

The procedure described in section **1** will be demonstrated by the following examples:

A) Let us consider a system with th action integral as

$$S(q_i) = \int dt \mathcal{L}(q_i, \dot{q}_i, t), \quad i = 1, ..., n, \tag{15}$$

where $\mathcal{L}$ is a regular Lagrangian with Hessian n. Parameterize the time $t \rightarrow \tau(t)$, with $\dot{\tau} = \frac{d\tau}{dt} > 0$. The velocities $\dot{q}_i$ may be expressed as

$$\dot{q}_i = q_i^{'} \dot{\tau}, \tag{16}$$

where $q_i^{'}$ are defined as

$$q_i^{'} = \frac{dq_i}{d\tau}. \tag{17}$$

Denote $t = q_0$ and $q_\mu = (q_0, q_i), \quad \mu = 0, 1, ..., n$, then the action integral (15) may be written as

$$S(q_\mu) = \int d\tau \dot{t} \mathcal{L}(q_\mu, \frac{q_i^{'}}{t}), \tag{18}$$

which is parameterization invariant since L is homogeneous of first degree in the velocities $q_\mu^{'}$ with L given as

$$L(q_\mu, \dot{q_\mu}) = \dot{t} \mathcal{L}(q_\mu, \frac{q_i^{'}}{t}). \tag{19}$$

The Lagrangian L is now singular since its Hessian is n.

The generalized momenta conjugated to the generalized coordinates q_μ are defined as

$$p^\tau_\mu = \frac{\partial L}{\partial q'_\mu}. \tag{20}$$

Therefore the i-th component is

$$p^\tau_i = \frac{\partial L}{\partial q'_i} = \frac{\partial \mathcal{L}}{\partial \dot{q}_i}, \tag{21}$$

and the zeroth component is

$$p_t = \frac{\partial L}{\partial \dot{t}} = \mathcal{L} - \frac{\partial \mathcal{L}}{\partial \dot{q}_i}\dot{q}_i, \tag{22}$$

Since $\mathcal{L}$ is regular, one can solve (21) for $\dot{q}_i$ in terms of terms of p_i and $\dot{q}_0$ as follows

$$\dot{q}_i = \dot{q}_i(p_i, \dot{q}_0) = w_i. \tag{23}$$

Substituting (23) in (22), one has

$$p_t = \mathcal{L}(q_i, \dot{q}_i(p_i, \dot{q}_0)) - p_i w_i. \tag{24}$$

The primary constraint is

$$H'_t = p_t + H_t = 0, \tag{25}$$

where H_t is defined as

$$H_t = -\mathcal{L}(q_i, w_i) + p_i w_i. \tag{26}$$

Calculation show that the canonical Hamiltonian

$$H_0 = -L(q_0, q_i, \dot{q}_0, w_i) + p^\tau_i q'_i + p_t \dot{q}_0 \mid_{pt=-H_t}, \tag{27}$$

vanishes identically.

The canonical method [1-4] leads us to obtain the set of Hamilton -Jacobi partial differential equations as follows:

$$H'_0 = 0, \tag{28}$$
$$H'_t = p_t + H_t = 0. \tag{29}$$

The equations of motion are obtained as total differential equations in many variables as follows:

$$dq^i = \frac{\partial H'_0}{\partial p_i} d\tau + \frac{\partial H'_t}{\partial p_i} dq^0 = \frac{\partial H'_t}{\partial p_i} dq^0, \tag{30}$$

$$dp^i = -\frac{\partial H'_0}{\partial q_i} d\tau + \frac{\partial H'_t}{\partial q_i} dq^0 = -\frac{\partial H'_t}{\partial q_i} dq^0, \tag{31}$$

$$dp_t = -\frac{\partial H'_0}{\partial q_0} d\tau + \frac{\partial H'_t}{\partial q_0} dq^0 = 0. \tag{32}$$

Since

$$dH'_t = dp_t + H_t, \tag{33}$$

vanishes identically, this system is integrable and the canonical phase space coordinates q_i and p_i are obtained in terms of the time ($q_0 = t$).

Till now we have discussed the Hamiltonian systems at the classical level. Quantization of system (15) can be achieved by means of path integral which we have discussed in reference [7], or the operator methods. We will deal with the later.

For Dirac quantization [5,6] of constrained dynamical systems, one takes the constraints equation as an operator whose action on the allowed Hilbert space vectors is constrained to $zero$, i., e., $H'_t \psi = 0$, we obtain

$$[\hat{p}_t + \hat{H}_t(\hat{q}_i, \hat{p}_i)]\psi = 0, \tag{34}$$

the ordinary Scrödinger equation. The resulting quantum theories are identical using either of the two-Lagrangians (15) or (16).

B) As a second example, let us consider a regular system with one-dimensional Lagrangian

$$l = \frac{1}{2}\dot{q}^2 - V(q), \tag{35}$$

where, $\dot{q} = \frac{dq}{dt}$. Replace the time by an arbitrary paremetrization $t = t(\tau)$ [10,13], we obtain the two dimensional Lagrangian

$$L = \dot{t}[\frac{1}{2}(\frac{q'}{\dot{t}})^2 - V(q)], \tag{36}$$

with $q' = \frac{dq}{d\tau}$ and $\dot{t} = \frac{dt}{d\tau}$. The Lagrangian L is singular since its Hessian is 1.

The generalized momenta $p_q^{(\tau)}$ and p_t are given by

$$p_q^{(\tau)} = \frac{\partial L}{\partial(\frac{\partial q}{\partial \tau})} = \frac{\partial l}{\partial \dot{q}} = p_q, \tag{37}$$

$$p_t = \frac{\partial L}{\partial(\frac{\partial t}{\partial \tau})} = -\frac{1}{2}(\frac{q'}{\dot{t}})^2 - V(q). \tag{38}$$

Since the rank of the Hessian is one we have one primary constraint as

$$H'_t = p_t + H_t = 0, \tag{39}$$

where H_t is defined as

$$H_t = \frac{p_q^2}{2} + V(q). \tag{40}$$

The canonical Hamiltonian which is obtained as

$$H = -L + p_q^{(\tau)} q' - H_t \dot{t}, \tag{41}$$

vanishes identically. We get the set of Hamilton-Jacobi partial differential equations as follows;

$$H'_t = p_t + \frac{p_q^2}{2} + V(q) = 0, \tag{42}$$

$$H' = 0 \tag{43}$$

The equations of motion are obtained as total differential differential equations as follows:

$$dq = \frac{\partial H'}{\partial p_q^{(\tau)}} d\tau + \frac{\partial H'_t}{\partial p_q^{(\tau)}} dt = p_q dt, \tag{44}$$

$$dp_q = -\frac{\partial H'}{\partial q} d\tau + \frac{\partial H'_t}{\partial q} dt = -\frac{dV}{dq} dt, \tag{45}$$

$$dp_t = -\frac{\partial H'}{\partial t} d\tau + \frac{\partial H'_t}{\partial t} dt = 0. \tag{46}$$

Since

$$dH'_t = dp_t + p_q dp_q + (\frac{dV}{dq}) dq, \tag{47}$$

vanishes identically, eqs. (44-46) are integrable and one obtains the dynamical variables q and p_q in terms of the physical time t.

Using the quantization procedure described in section **1**, we obtain

$$[\hat{p_t} + \frac{1}{2}(\hat{p_q})^2 + V(q)]\psi = 0, \tag{48}$$

which is the Schrödinger equation for linear harmonic oscillator.

3 Relativistic particle

In this section we shall use the canonical method [1-4] to obtain the equations of motion for a relativistic particle of charge e in an external electromagneticthis system and for a free relativistic particle.

A)Let us consider the action for a relativistic charged particle in an external field as

$$S = \int L d\tau, \tag{49}$$

where the Lagrangian L is given by

$$L = -[mc\sqrt{g^{\mu\nu}\dot{q_\mu}\dot{q_\nu}} + \frac{e}{c}\dot{q_\mu}A^\mu], \quad \mu, \nu = 0, 1, 2, 3, \tag{50}$$

and $g^{\mu\nu} = diag(1, -1, -1, -1)$. The momenta conjugated to $\dot{q_\mu}$ are

$$p_\mu = \frac{\partial L}{\partial(\dot{q^\mu})} = -(\frac{mc\dot{q_\mu}}{\sqrt{\dot{q}^2}} + \frac{e}{c}A_\mu). \tag{51}$$

Hence, the zeroth component p_0 and the ith component p_i are given as

$$p_0 = -(\frac{mc\dot{q_0}}{\sqrt{\dot{q}^2}} + \frac{e}{c}A_0), \tag{52}$$

$$p_i = -(\frac{mc\dot{q_i}}{\sqrt{\dot{q}^2}} + \frac{e}{c}A_i). \tag{53}$$

Since the rank of the Hessian matrix is three, one can solve (53) to obtain the velocities $\dot{q_i}$ in terms of $\dot{q_0}, p_i$ and A_i as

$$\dot{q_i} = \frac{k_i\dot{q_0}}{[|\mathbf{k_i}|^2 + m^2c^2]^{1/2}} = w_i, \tag{54}$$

where k_i is defined as

$$k_i = (p_i + \frac{e}{c}A_i). \tag{55}$$

Substituting (54) in (52) one gets

$$p_0 = -[|\mathbf{k_i}|^2 + m^2c^2]^{1/2} - \frac{e}{c}A_0 = -H_0. \tag{56}$$

Now the primary constraint is given as

$$H_0' = p_0 + [|\mathbf{k_i}|^2 + m^2c^2]^{1/2} + \frac{e}{c}A_0 = 0. \tag{57}$$

The canonical Hamiltonian (3) can be written as

$$H = -L + p_i w_i + p_0 \dot{q}_0. \tag{58}$$

Explicit calculations show that H vanishes identically and lead to

$$H' = 0. \tag{59}$$

Equations (57) and (59) lead us to obtain the set of Hamilton- Jacobi partial differential equations as follows

$$H' = 0, \tag{60}$$

$$H_0' = p_0 + H_0 = 0. \tag{61}$$

The equations of motion are obtained as total differential equations as follows

$$dq_i = \frac{\partial H'}{\partial p_i}d\tau + \frac{\partial H_0'}{\partial p_i}dq_0, \tag{62}$$

$$dq_i = \frac{k_i}{[|\mathbf{k_i}|^2 + m^2c^2]^{1/2}}dq_0, \tag{63}$$

$$dp_i = -\frac{\partial H'}{\partial q_i}d\tau - \frac{\partial H_0'}{\partial q_i}dq_0, \tag{64}$$

$$dp_i = -\frac{e}{c}[\frac{\partial A_j}{\partial q^i}\frac{k_i}{[|\mathbf{k_i}|^2 + m^2c^2]^{1/2}} + \frac{\partial A_0}{\partial q^i}]dq_0, \tag{65}$$

$$dp_0 = -\frac{\partial H'}{\partial q_i}d\tau - \frac{\partial H_0'}{\partial q_0}dq_0, \tag{66}$$

$$dp_0 = -\frac{e}{c}[\frac{\partial A_j}{\partial q^0}\frac{k_i}{[|\mathbf{k_i}|^2 + m^2c^2]^{1/2}} + \frac{\partial A_0}{\partial q^0}]dq_0. \tag{67}$$

In order to have a consistent theory, one should consider the total variation of H' and H'_0. Since H' vanishes identically, we shall consider only the total variation of H'_0. In fact

$$dH'_0 = \frac{k_i}{[|\mathbf{k_i}|^2 + m^2c^2]^{1/2}}(dp_i + \frac{e}{c}dA_i) + dp_0 + \frac{e}{c}dA_0. \tag{68}$$

Making use of equations (62-67), one gets

$$dH'_0 = 0. \tag{69}$$

Since the variations of H' and H'_0 are identically zero, no further constraints arise. Hence, the equations of motion are integrable and the canonical phase space coordinates are obtained in terms of parameter q_0.

We can use the quantization procedure discussed in previous sections to obtain

$$[\hat{p}_0 + \sqrt{(\hat{p}_i + \frac{e}{c}\hat{A}_i)^2 + m^2c^2} + \frac{e}{c}\hat{A}_0]\psi = 0. \tag{70}$$

B) To obtain the equations of motion for a free relativistic particle we can use the results obtained in section A, but instead we consider no interaction with external field ($A_\mu = 0$). In this case the equations of motion are obtained as

$$dq_i = \frac{\partial H'}{\partial p_i}d\tau + \frac{\partial H'_0}{\partial p_i}dq_0, \tag{71}$$

$$dq_i = \frac{p_i}{[|\overrightarrow{p}|^2 + m^2c^2]^{1/2}}dq_0, \tag{72}$$

$$dp_i = -\frac{\partial H'}{\partial q_i}d\tau - \frac{\partial H'_0}{\partial q_i}dq_0, \tag{73}$$

$$dp_i = 0, \tag{74}$$

$$dp_0 = -\frac{\partial H'}{\partial q_i}d\tau - \frac{\partial H'_0}{\partial q_0}dq_0, \tag{75}$$

$$dp_0 = 0. \tag{76}$$

Equations (71-76) are integrable [7]. Hence, we obtain

$$p_i = a_i, \tag{77}$$

and

$$p_0 = c_i, \tag{78}$$

where $a_i$and c_i are arbitrary constants. So integration of (72) will give

$$q_i = \frac{p_i}{[|\overrightarrow{p}|^2 + m^2c^2]^{1/2}} q_0 + \alpha_i, \tag{79}$$

where α_i are constants.

Again as for the relativistic charged particle we can quantize this system to obtain

$$[\hat{p}_0 + \sqrt{(\hat{\overrightarrow{p}} + m^2c^2}]\psi = 0. \tag{80}$$

4 Hamilton-Jacobi treatment of the field systems

The Hamilton-Jacobi treatment for finite dimensional systems which are discussed in previous sections will be extended to the field theory. Let us consider the action of this theory as

$$S[\phi_A] = \int d^{D+1}t\mathcal{L}(\phi_A, \partial_a\phi_A), \quad a = 0, 1, ..., D, \quad A = 1, ..., N. \tag{81}$$

where $\mathcal{L}$ is a regular Lagrangian density. Introduce $D+1$ parameters T^a, with $det \mid \frac{\partial T^a}{\partial t^b} \mid > 0$, and treat the t^a on the same footing as the original field ϕ_A. The action (81) is now reads as

$$S[\phi_A, t^a] = \int d^{D+1}T\mathcal{L}_T(\phi_A, \partial_a\phi_A), \tag{82}$$

where

$$\mathcal{L}_T = \mathcal{L}(\phi_A, \frac{\partial\phi_A}{\partial T^a}(J^{(-1)})^b_a) \det(J^b_a), \tag{83}$$

$$J^b_a = \frac{\partial t^a}{\partial T^b}. \tag{84}$$

The action (82) is invariant under parameterization $T \rightarrow T'(T)$. Now the Lagrangian $\mathcal{L}_T$ is singular since it is linear in the velocities $\frac{\partial t^a}{\partial T^0}$. The momenta are now

$$\pi^A = \frac{\partial \mathcal{L}_T}{\partial(\frac{\partial \phi_A}{\partial T^0})} = \frac{\partial \mathcal{L}}{\partial(\frac{\partial \phi_A}{\partial t^0})}, \tag{85}$$

$$\pi^a = \frac{\partial \mathcal{L}_T}{\partial(\frac{\partial t^a}{\partial T^0})}. \tag{86}$$

Since $\mathcal{L}$ is regular, one can solve (85) for $\frac{\partial \phi_A}{\partial T^0}$ in terms of π^A, ϕ_A, t_a and $\frac{\partial t^a}{\partial T^0}$. Substituting (85) in (86), one has

$$\pi_a = -\frac{\partial \mathcal{L}}{\partial(\partial_a \phi_A)} \partial_a \phi_A + \delta_a^b \mathcal{L}. \tag{87}$$

Hence, the primary constraint is

$$H'_a = \pi_a + H_a(\pi_A, t^a, \phi^A) = 0, \tag{88}$$

where

$$H_a = (\frac{\partial \mathcal{L}}{\partial(\partial_a \phi_A)} \partial_a \phi_A - \delta_a^b \mathcal{L}) Adj(J_{b0}). \tag{89}$$

Besides the canonical Hamiltonian H_T is defined as

$$H_T = \int d^D T (\pi^A \frac{\partial \phi_A}{\partial T^0} + \pi^a \frac{\partial t^a}{\partial T^0}|_{\pi_a = -H_a} - \mathcal{L}_T). \tag{90}$$

Calculations show that H_T vanishes identically.

Now we would like to obtain the equations of motion for this system using the canonical method [1-4]. Making use of equations (88) and (90), one obtains the set of Hamilton-Jacobi partial differential equations as

$$H'_T = 0, \tag{91}$$

$$H'_a = \pi_a + H_a = 0. \tag{92}$$

The equations of motion are obtained as total differential equations as follows:

$$d\phi^A = \frac{\partial H'_T}{\partial \pi_A} dT^a + \frac{\partial H'_a}{\partial \pi_A} dt^a = \frac{\partial H'_a}{\partial \pi_A} dt^a, \tag{93}$$

$$d\pi^A = -\frac{\partial H'_T}{\partial \phi_A} dT^a - \frac{\partial H'_a}{\partial \phi_A} dt^a = -\frac{\partial H'_a}{\partial \phi_A} dt^a, \tag{94}$$

$$d\pi_a = -\frac{\partial H'_T}{\partial t^a} dT^a - \frac{\partial H'_a}{\partial t^a} dt^a = \frac{\partial H'_a}{\partial t^a} dt^a. \tag{95}$$

Since

$$dH'_a = d\pi_a + dH_a, \tag{96}$$

vanishes identically, the set of equations (93-95) is integrable and the canonical phase space coordinates ϕ^A and π^A are obtained in terms of t^a.

To obtain the path integral quantization for this system, we can use the canonical path integral formulation discussed in references [7-10]. Making use of (7), the canonical action integral is calculated as

$$S[\phi^A] = \int [-H_a + \pi^A \frac{\partial H'_a}{\partial \pi_A}] dt^a. \tag{97}$$

Then the path integral integral for the field system is given as

$$\langle Out \mid S \mid In \rangle = \int \prod_{A=1}^{N} d\phi^A \; D\pi^A \exp i \int_{t^a}^{t'^a} [-H_a + \pi^A \frac{\partial H'_a}{\partial \pi_A}] dt^a. \tag{98}$$

Integrating over π^A gives

$$\langle Out \mid S \mid In \rangle = \int \prod_{A=1}^{N} d\phi^A \exp i \{ \int_{t^a}^{t'^a} d^{D+1}t \; \mathcal{L}(\phi_A, \partial_a \phi_A) \}. \tag{99}$$

5 Conclusion

In this work we use the canonical method to investigate the significance of time for constrained systems with weakly vanishing Hamiltonian. For these systems we obtain the dynamical variables in terms of physical time t without fixing any gauge.

In general, the determination of the Hamiltonians H'_α is the crucial step. If the system is integrable (Jacobi system), then the equations of motion are obtained in terms of t_α.

for the first and the second example, the equations of motion are obtained in terms of two parameters τ and t. However, since the Hamiltonian H' vanishes identically, the parameter τ doe not appear in the equations. Hence, any parameter can be replaced by τ and one can use t as evolution parameter without losing the Hamiltonian structure of the equations of motion.

Besides, as for the first two examples, without fixing any gauge, the equations of motion for a relativistic particle discussed in section **3** are obtained in terms of parameter q^0.

In section **4** we have obtained the equations of motion for field systems using the Hamilton-Jacobi method. In general, continuous systems (fields) are treated as systems with infinite degrees of freedom. The Lagrangian density is defined and the usual variational principle technique is used to obtain the equations of motion. the quantities $t^a = x, y, z, t$ are completely independent indices of the theory. However, using the parameterization $t^a \rightarrow T^a$ to these theories leads to obtain the equations of motion in terms of t^a and T^a. Moreover, since the Hamiltonian H'_T vanishes identically, the canonical phase space coordinates ϕ^A and π^A are obtained in terms of t^a and the parameters T^a do not appear in the equations of motion. On the other hand the path integral for this system is obtained as an integration over the canonical phase space coordinates ϕ^A and π^A without using any gauge fixing conditions.

As a conclusion it is obvious that the canonical method leads to obtain the dynamical variables for finite dimensional systems with weakly vanishing Hamiltonian in terms of the time t without using any gauge fixing conditions. The extension of this treatment to field systems leads to obtain the equations of motion in terms of parameters t^a and the number of independent parameters depend on the Hessian and the integrability conditions.

The operator quantization for the above system is obtained using the Hamilton Jacobi method without using any gauge fixing conditions. In this case the evolution of the state Ψ is described by the Schrödinger equations $(i\frac{\partial \Psi}{\partial t} = \hat{H}\Psi)$ (34, 48, 70, 80). Where the operators $\hat{H}$ are the quantum Hamiltonians corresponding to the classical physical Hamiltonians given in equations (26, 40, 56, 80).

References

[1] Y. Güler, Nuovo Cimento B, 107 (1992)1389.

[2] Y. Güler, Nuovo Cimento B, 107 (1992)1143.

[3] S. I. Muslih and Y. Güler, Nuovo Cimento B, 110 (1995) 307.

[4] S. I. Muslih and Y. Güler, Nuovo Cimento B, 113 (1998)277.

[5] P. A. M. Dirac, *Lectures on Quantum Mechanics*,Belfer Graduate School of Science, Yehiva University (A cademic Press, New York)1964.

[6] P. A. M. Dirac, Can. J. Math., 2 (1950)129.

[7] S. I. Muslih, Nuovo Cimento B, 115 (2000)1.

[8] S. I. Muslih, Nuovo Cimento B, 115 (2000)7.

[9] S. I. Muslih," Path integral quantization of Yang-Mills theory" submitted fo Nuovo Cimento B.

[10] S. I. Muslih, Hadronic J, 23 (2000)203.

[11] A. Hanson, T. Regge and C. Teitelboim, *Constrained Hamiltonian Systems* (Accademia Nazionole dei Lincei, Rome) 1976.

[12] D. M. Gitman and I. Tyutin, *Quantization of Fields with Constraints* (Springer- Verlag, Berlin, Heidelberg) 1990.

[13] J. Goldeberg, E. Newman and C. Rovelli, J. Math. Phys. 32 (1991) 2739.